

Possible bipolar global expression of the P3 and P4 glacial events of eastern Australia in the Northern Hemisphere: Marine diamictites and glendonites from the middle to upper Permian in southern Verkhoyanie, Siberia

V.I. Davydov^{1,2,*}, R.V. Kutugin³, and E.V. Karasev⁴

¹Department of Geosciences, Boise State University, 1910 University Drive, Boise, Idaho 83725, USA

²Institute of Petroleum Geology and Geophysics, Prospekt Ac. Koptjugy 3, Novosibirsk 630090, Russia

³Diamond and Precious Metal Geology Institute, Siberian Branch, Russian Academy of Sciences, 39 Lenina Prospect, Yakutsk 677891, Russia

⁴Paleontological Institute RAS, Profsovnaya 123, Moscow 117647, Russia

We are pleased to receive the Comment of John Isbell (Isbell, 2023), who is a leading specialist in the Late Paleozoic glacial studies. We have to agree with Isbell that the origin of the Siberian sediments of interest is complicated and not fully documented or understood (Davydov et al., 2022). The region is so large, and currently so remote, that we were not able to study all the necessary aspects of glacial-marine sediments, especially laterally toward the areas where the real continental glaciation might have developed.

In our paper, we proposed that the middle-late Permian glacial events were expressed as glacial-marine sediments originating from perennial oceanic and coastal sea ice, similar to the setting in the present-day Arctic Ocean. The recent investigations of the evolution and extent of sea ice in modern seas and oceans demonstrate that sea ice appears only after the formation of a large ice sheet (DeConto et al., 2007; Gowan et al., 2021). Therefore, in the present-day Arctic Ocean, perennial oceanic and coastal sea ice along the northern coast of Siberia is a derivative of the Greenland Ice Sheet (Törnqvist and Hijma, 2012; Vernal et al. 2020).

The Southern Verkhoyanie foredeep, during the middle-late Permian, was bordered to the northeast by the Siberian continent, and glacial ice sheets potentially could have developed in the highlands and mountains in the north and northeast of the continent and in the adjacent Argun (Stanovoy) terrane (Parfenov et al., 2011) (Fig. 1). The occurrence of the yet-undiscovered ice sheet would have facilitated the development of sea ice in the areas surrounding the ice sheet and along the western margin of the Siberian continent (Fig. 1). During Late Paleozoic, the Southern Verkhoyanie and adjacent areas, including the Okhotsk-Taigonos Arc, the Okhotsk Massif, and the northern margin of the Siberian Platform (recent southern and southeast Siberia), were the only areas in the Northern Hemisphere where glacial deposits could have developed (Fig. 1). Our previous study in the Aiyun-Yuryak and Priokhotie regions (Isbell et al., 2016) suggested that the Capitanian volcanoclastic sediments of the Atkan Formation (widely distributed in northeastern Russia) are turbidites deposited as prograding and abandoning sediment gravity-flow fans, with chaotic and folded strata having formed as slumps. In our current study, the Capitanian rocks from the Kobume River were interpreted as an ice-free area (Davydov et al., 2022). We are expecting to soon obtain radioisotopic ages from the glacial-marine sediments in Kobume, which might clarify the model and the chronostratigraphic constraints on the glacial episodes in this intriguing, but still very poorly studied region.

ACKNOWLEDGMENTS

This study is a part of the Ministry of Science and Higher Education of Russia project FWZZ-2022-0005 (to V. Davydov) performed by the Institute of Petroleum Geology and Geophysics, Novosibirsk, and FUEM-2019-0002 (to R. Kutugin) performed by the DPMGI SB RAS.

REFERENCES CITED

- Davydov, V., Budnikov, I., Kutugin, R., Nurgalieva, N., Biakov, A., Karasev, E., Kilyasov, A., and Makoshin, V., 2022, Possible bipolar global expression of the P3 and P4 glacial events of eastern Australia in the Northern Hemisphere: Marine diamictites and glendonites from the middle to upper Permian in southern Verkhoyanie, Siberia: *Geology*, v. 50, p. 874–879, <https://doi.org/10.1130/G50165.1>.
- DeConto, R.; Pollard, D.; Harwood, D., 2007, Sea ice feedback and Cenozoic evolution of Antarctic climate and ice sheets: *Paleoceanography*, 22 (3), 1–18. DOI: 10.1029/2006PA001350.
- Gowan, E. J.; Zhang, Xu; Khosravi, S.; Rovere, A.; Stocchi, P.; Hughes, A. L. C. et al., 2021, A new global ice sheet reconstruction for the past 80 000 years: *Nature communications* 12 (1), p. 1199. DOI: 10.1038/s41467-021-21469-w.
- Isbell, J.L., Biakov, A.S., Vedernikov, I.L., Davydov, V.I., Gulbranson, E.L., and Fedorchuk, N.D., 2016, Permian diamictites in Northeastern Asia: Their significance concerning the bipolarity of the late Paleozoic ice age: *Earth-Science Reviews*, v. 154, p. 279–300, <https://doi.org/10.1016/j.earscirev.2016.01.007>.
- Isbell, J.L., 2023, Possible bipolar global expression of the P3 and P4 glacial events of eastern Australia in the Northern Hemisphere: Marine diamictites and glendonites from the middle to upper Permian in southern Verkhoyanie, Siberia: *Comment: Geology*, v. 51, p. e558. <https://doi.org/10.1130/G51125C.1>.
- Parfenov, L.M., Nokleberg, W.J., Berzin, Nikolai A., Badarch, G., Dril, S.I., Gerel, O., et al., 2011, Tectonic and metallogenic model for northeast Asia: US Geological Survey Open-File Report.
- Törnqvist, T. E., Hijma, M. P. (2012). Links between early Holocene ice-sheet decay, sea-level rise and abrupt climate change: *Nature Geosciences* 5 (9), pp. 601–606. DOI: 10.1038/ngeo1536.
- Vernal, A. de, Hillaire-Marcel, C., Le Duc, C., Roberge, Ph., Brice, C., Matthiessen, J. et al.: 2020, Natural variability of the Arctic Ocean sea ice during the present interglacial: *Proceedings of the National Academy of Sciences of the United States of America* 117 (42), pp. 26069–26075. DOI: 10.1073/pnas.2008996117.




Figure 1. Potential distribution of the ice sheet and sea-ice in the northern Hemisphere compared with the glacial development in the southern Hemisphere (middle-late Permian).